## **TBI: Mortality and Nonfatal Hospitalization**

## **Data Analysis Report**

Minnesota 2006

**Prepared by:** 

Anna Gaichas, MS Leslie Seymour, MD MPH Jon Roesler, MS Ayo Adeniyi, MPH MBBS Mark Kinde, MPH

Minnesota Department of Health St. Paul, Minnesota

For the

**Centers for Disease Control and Prevention** 

As part of

TBI Extended Surveillance Program (TBIES) / PA 05027, Award Number 524800-01

October 30, 2008

### **Table of Contents**

Text Overview1	
Table 1: TBI Mortality Number and Rate by Gender, Age Group, and Leading Causes, Minnesota 2006	2
Figure 1: TBI Mortality Rate by Year, Minnesota 1998-2006.	;
Table 2: Nonfatal Hospitalized TBI Number and Rate by Gender, Age Group, and Leading Causes, Minnesota 20064	1
Figure 2: Nonfatal Hospitalized TBI Rate by Year, Minnesota 1993-2006	i

Suggested citation: Gaichas A, Seymour L, Roesler J, Adeniyi A, Kinde M. *TBI: Mortality and Nonfatal Hospitalization Data Analysis Report, Minnesota 2006.* St. Paul: Minnesota Department of Health, October 2008.

#### **Text Overview**

**Background:** Minnesota is one of only four states funded by the CDC to conduct surveillance of Emergency Department (ED)-treated traumatic brain injury (TBI) for the 2005-2010 data period.

**Methods:** Electronic, centralized discharge data from death certificates, the Minnesota TBI/Trauma Registry and the Minnesota Hospital Association were analyzed using the CDC's TBI case definition. Analysis for this report was limited to Minnesota residents.

**Results:** In 2006, there were 856 TBI deaths, an adjusted rate of 16.0 per 100,000 population. This rate is similar to those of the previous three years. The number of nonfatal hospitalized (NH) TBIs in 2006 was 4,867, yielding an adjusted rate of 92.8 per 100,000 population. The data show an increasing trend. Comparison with 1993-1997 is difficult because of changes in case identification methods.

The TBI mortality data show a bi-modal curve, with peaks among adolescents/young adults and the elderly. The TBI-NH data show a tri-modal curve, with peaks among the infants, adolescents/young adults, and the elderly. Both the TBI mortality and TBI-NH peaks occur in older adults, with the highest rates among persons age 85 and above (101.10 per 100,000 population for mortality and 454.3 per 100,000 population for TBI-NH). The TBI mortality and TBI-NH rates for males were consistently higher than those for females, with male rates among infants and adolescents/young adults approximately twice that of the female rates. Male TBI mortality rates were approximately three times that of the female rates for elders 80-84 and male TBI-NH rates were about three times higher than female rates for young adults age 20-29. Some age- and gender-specific TBI mortality rates were not calculated due to small numbers.

The leading causes of TBI mortality were firearms, especially self-inflicted, unintentional falls, and motor vehicle traffic (MVT). The leading causes of TBI-NH were unintentional falls, MVT, and assault. TBI deaths from firearms were concentrated around middle-aged adults, while MVT fatalities were most prevalent in teens and young adults and unintentional fall deaths most often occurred in older adults. For TBI-NH, unintentional fall rates showed a bi-modal curve with the highest rates among older adults 65 years and above with a smaller peak in infants age < 1. MVT and assault TBI-NH rates were highest for adolescents/young adults, ages 15-24 for MVT and ages 20-29 for assault.

**Discussion:** Data in this report show the importance of TBI as a public health problem. Population-based information regarding TBI deaths and hospitalizations can be useful in assessing the effect of prevention efforts. Additionally, abstracted data will yield important information on demographic features (e.g., race/ethnicity), modifiable risk factors (e.g., alcohol use), and circumstances at time of injury (e.g., specific activity in which the individual was engaged at time of injury) that in turn can be used to guide intervention activities.

		All caus	ses		Firearm				l,	Motor vehicle traffic			
		total	-	Total		Self-inf	licted	uninten	tional	Tota		Occu	pant
Total		N	Rate	Ν	Rate	Ν	Rate	N	Rate	Ν	Rate	N	Rat
	All Ages	819	15.4	223	4.3	193	3.7	241	4.4	223	4.2	138	2.
	<1	6		0		0		0		3		3	
	1-4	12 z		0	•	0 0		2 0		1 a		0	
	5-9	0 12		U 3		0		0		2 9		0	
	10-14	50	13.4	10		7		1		33	8.9	26	7.0
	20-24	58	15.5	18		11	-	1		34	9.1	16	
	25-29	51	14.5	27	7.7	20	5.7	1		17		11	
	30-34	36	11.1	13		11	•	2	•	16	•	12	
	35-39	43	12.0	17		15	. 57	7		6 14	•	2	
	40-44	50 67	12.3	24 25	6.0	25 21	5.7 5.0	5 13		14 16	•	5 10	
	43-49	53	14.3		6.2	21	5.7	10		10		10	
	55-59	63	20.1	17		16		16		17		10	
	60-64	30	13.5	6		6	•	12		4	•	3	
	65-69	30	17.5	9	•	9		5		11		7	
	70-74	39	28.2	8	•	8		23	16.6	6	•	4	
	75-79	53 67	44.1 70.1	13	•	13		29 43	24.1 45.0	0 10	•	5	
	00-04 85±	93	91.5	, 3		3			69.9	10		5	
Male													
	All Ages	590	23.7	203	7.9	182	7.2	135	5.9	150	5.8	88	3.4
	<1	4		0		0		0		1		1	
	1-4	9	•	0	•	0	•	1		1	•	0	
	5-9	1		0		0		0		1		0	
	10-14	34	179	3 8	•	2		1		4 20	10 5	17	
	15-19 20-24	45	23.6	16		11		0		20	12.6	10	
	25-29	43	23.6	24	13.2	19		1		12		7	
	30-34	31	18.8	13	•	11	•	2	•	12	•	10	
	35-39	36	19.8	15		13		5		5		1	
	40-44	43 52	21.3	23 20	11.4	22 19	10.9	5 0		10 12	•	1 0	
	45-49	33 46	25.1 24.7	20 21	11.3	21	11.3	2		8		0 7	
	55-59	48	30.7	16		16		13		12		6	
	60-64	24	22.0	5		5		9		4		3	
	65-69	24	29.2	8		8		4		7		5	
	70-74	28	43.9	8	•	8	•	15	•	4	•	2	
	75-79	39 36	/4.6 06.2	13	•	13	•	16 18	•	5	•	4	
	00-04 85⊥	39	125.6	, 3	•	3		28	90.2	2		0	
Female													
	All Ages	229	7.7	20	0.7	11		106	3.2	73	2.7	50	1.8
	<1	2		0		0		0		2		2	
	1-4	3		0		0		1		0	•	0	
	5-9	2		U		0		0		1 5		U 4	
	10-14 15-10	16	•	2	•	2	•	0	•	13	•	9	
	20-24	13		2		0		1		10		6	
	25-29	8		3		1		0		5		4	
	30-34	5	•	0	•	0	•	0	•	4	•	2	
	35-39	7	•	2	•	2		2		1	•	1	
	40-44 45 40	/ 14		1 5		1 2		U 21	2000-000-000-000-000-000-000-000-000-00	4 3		2	
	+3-+7 50-54	* <b>*</b> * 7		2		0		2		3		3	
	55-59	15		1		0		3		5		4	
	60-64	6	•	1	•	1	•	3	•	0	•	0	
	65-69	6		1	•	1		1		4		2	
	70-74	11	•	0	•	0	•	8	•	2	•	2	
	75-79	14 31	53 2	0		0	•	13	43.0	1	•	1	
	851 851	54	55.5 76 5	U N		n N	•	2.3 43	40.0 60.9	י 5		1 5	
	UN T					<u>aanaa dadda Mali</u>			na se	nn an tha	ooraadde fiddif	non de la Ciè	

# Figure 1: TBI Mortality Rate by Year Minnesota 1998-2006



Rates are adjusted to the U.S. 2000 standard population.

Table 2	le 2: Nonfatal Hospitalized TBI Number and Rate by Gender, Age Group, and Leading Causes, Minnesota 2 Motor vehicle traffic Assault												ota 2006	
			11565	Fa	n			mere ti ai	inc		Assault Struck by or			
			al	uninter	ntional	Тс	otal	Occ	unant	Тс	otal	aga	ninst	
Total		N	Rate	N	Rate	N	Rate	- N	Rate	N	Rate		Rate	
-	All Ages	4,662	88.6	2,039	38.3	1,334	25.6	944	1 18.1	373	7.1	248	3 4.7	
	<1	87	118.3	57	77.5	4	· .	. 4	ι.	12	2.	. (	).	
	1-4	111	40.9	67 - 1	24.7	16		13	2000-000-000-000-000-00-00-00-00-00-00-0	5		. (	) . 2000-000-000-000-000-000-000-000-000-00	
	5-9 10-14	135 174	40.5 49.4	51 44	15.3	39 46	11./	15	) 57	1	)		) )	
	15-14	470	126.1	51	13.7	247	66.3	206	5 55.3	45	12.1	. 32	, . 2. 8.6	
	20-24	395	105.6	43	11.5	187	50.0	147	7 39.3	76	5 20.3	58	15.5	
	25-29	281	80.0	47	13.4	126	35.9	91	25.9	55	15.7	36	i 10.2	
	30-34	199	61.2	26	8.0	96	29.5	69	) 21.2	36	5 11.1	20	) 6.2	
	35-39 40-44	230	04.2 70.1	88	22.0	80 96	22.3	54 0 6 <sup>2</sup>	2 14.5 3 15.7	32 27	2 8.9 2 6.7	23	) 7.0	
	45-49	273	64.9	86	20.5	92	21.9	52	2 12.4		8.8	2.	6.0	
	50-54	236	63.7	94	25.4	70	18.9	38	3 10.3	27	7.3	17	· ·	
	55-59	223	71.1	112	35.7	65	20.7	39	) 12.4	7		. 7		
	60-64 65 60	172	77.4	98 195	44.1	42 25	18.9	) 28 ac	3 12.6	5	) 		) 	
	03-09 70-74	16 <del>9</del> 187	135.3	12.3	74.0 99.1	26 26	18.8	20	10.5	1	19999999999999999999999999999999999999	. (	)	
	75-79	261	217.0	211	175.4	23	19.1	20	) 16.6	1		. 1		
	80-84	300	313.9	239	250.1	28	29.3	21	1 22.0	1		. 1		
	85+	458	450.6	402	395.5	16		. 14	l.	1		. (	)	
Male	A 11 A	2.020	117.1	N	Kate	N		57(		202				
	All Ages	2,929	117.1	1,115	47.2 88.0	205	33.0	5/(	) 21.8	502 8	2 11.3 R	212	) 7.9	
	1-4	60	43.3	30	21.6	10			3.	2	· · ·	. (	).	
	5-9	82	48.1	32	18.8	21	12.3	ç	).	1			) .	
	10-14	115	63.6	32	17.7	29	16.0	1(	) .	0	) .	. (	) .	
	15-19	325	171.0	37	19.5	151	79.4	118	3 62.1	41	21.6	) 29 5/	15.3	
	20-24	213	116.7	30 37	20.3	139 94	51.5	60	37.8	42	23.0	29	· 20.3	
	30-34	139	84.2	14	•	63	38.2	4(	) 24.2	29	) 17.6	17		
	35-39	165	90.9	42	23.1	51	28.1	29	9 16.0	27	14.9	22	. 12.1	
	40-44	187	92.8	64	31.8	55	27.3	36	5 17.9	21	10.4	. 15		
	45-49	188	88.9 83.2	02 59	29.3	04 46	29.3	20	5 15.2 I 113	20 20	) 12.3	14 14	lininininini L	
	55-59	150	95.8	71	45.4	44	28.1	23	3 14.7	6	j 10.7	. 6		
	60-64	106	97.4	58	53.3	25	23.0	13	3.	5	;		ί.	
	65-69	113	137.3	73	88.7	19		. 12	2	4				
	70-74 75-70	103	161.4	/0 108	109.7	1/		. 12	· ·	. ()	) .	. (	) .	
	80-84	140	417.0	100	326.1	13		. 11	- 	1		. 1		
	85+	168	540.9	141	454.0	7		. (	5	1			) .	
Female		Ν	Rate	N	Rate	N	Rate	e N	N Rate	e N	Rate	e N	Rate	
	All Ages	1,733	61.5	924	30.5	469	18.1	374	4 14.4	71	2.7	36	5 1.4	
	<1 1 4	34	94.4 28.2	24	66.6	2		. 2	2.	. 4	,	. (	).	
	1-4 5-9	53	30.3 32.6	37 19	21.0	18		1	) )	(	)	. (	, )	
	10-14	59	34.5	12	•	17	•	. 1(	) .	0	) .	. (	) .	
	15-19	145	79.4	14		96	52.6	88	3 48.2	4	Ŀ.	. 3	з.	
	20-24	84 69	45.8	13		48	26.2	36	5 19.6	9	) 	. 4		
	25-29 30-34	08 60	40.5 37.5	10 12	1000000000K	94 33	18.9 20.7	<u>4</u> 20	4 13.0 ) 18.1	13 7	1		100000000000000 1	
	35-39	65	36.8	12		29	16.4	23	3 13.0	5	5		· ·	
	40-44	94	47.1	24	12.0	41	20.6	27	7 13.5	6	5	. 4	ŀ .	
	45-49	85	40.7	24	11.5	30	14.4	. 24	4 11.5	11				
	50-54 55-59	81 72	43.9	35	19.0	24	13.0	17	/ . 5	. 7	· .	. 3	· .	
	55-59 60-64	73 66	40.5 58.3	41	20.1 35.3	21 17	15.4	· 10	5.		) .		) .	
	65-69	76	85.0	52	58.2	16		10	5	Õ	)	. Č		
	70-74	84	113.0	67	90.1	9		. 8	3.	1	•	. (	).	
	75-79	121	178.0	103	151.6	10		. 8	3.	0	).	. (	).	
	δυ-δ4 95	144 200	247.6	117 041	201.2	11		. 1(	ן. א	. 0	)	. (		

Rates are per 100,000 populati	on. Total rates are adjusted to the U.S	. 2000 standard population.
--------------------------------	---	-----------------------------

# Figure 2: Nonfatal Hospitalized TBI Rate by Year Minnesota 1993-2006



Rates are adjusted to the U.S. 2000 standard population.